

AD705164

FOREIGN TECHNOLOGY DIVISION



COMPARATIVE CHARACTERISTIC OF SOME OF THE DEVICES USED
TO DETERMINE THE MICROBIAL CONTAMINATION OF THE AIR

by

G. N. Ishchenko, K. Khamrakulova, and R. Samigullin



Distribution of this document is unlimited. It may be released to the Clearinghouse, Department of Commerce, for sale to the general public.

Reproduced by the
CLEARINGHOUSE
for Federal Scientific & Technical
Information Springfield Va. 22151

EDITED TRANSLATION

COMPARATIVE CHARACTERISTIC OF SOME OF
THE DEVICES USED TO DETERMINE THE MICROBIAL
CONTAMINATION OF THE AIR

By: G. N. Ishchenko, K. Khamrakulova, and
R. Samigullin

English pages: 4

Source: Meditsinskiy Zhurnal Uzbekistana
(Uzbekistan Journal of Medicine),
No. 6, 1962, pp. 16-18.

Translated by: V. Mesenzeff/TDBRS-3

UR/0242/62/000/006

THIS TRANSLATION IS A RENDITION OF THE ORIGINAL FOREIGN TEXT WITHOUT ANY ANALYTICAL OR EDITORIAL COMMENT. STATEMENTS OR THEORIES ADVOCATED OR IMPLIED ARE THOSE OF THE SOURCE AND DO NOT NECESSARILY REFLECT THE POSITION OR OPINION OF THE FOREIGN TECHNOLOGY DIVISION.

PREPARED BY:

TRANSLATION DIVISION
FOREIGN TECHNOLOGY DIVISION
WP-AFB, OHIO.

COMPARATIVE CHARACTERISTIC OF SOME OF THE DEVICES USED
TO DETERMINE THE MICROBIAL CONTAMINATION OF THE AIR

G. N. Ishchenko, K. Khamrakulova, and R. Samigullin

Department of Microbiology
(Head — Professor F. I. Shevchenko)
Samarkand Medical Institute

To study microbial contamination of the air, several special devices — bacteria traps — are recommended. However, they do not fully satisfy the requirements of practicing laboratories.

We directed our attention to the fact that when air is seeded with Krotov apparatus the indices of microbial contamination are considerably lower than those from other methods and devices. This prompted us to carry out a comparative study of the results obtained with Krotov and P. P. D'yakonov apparatus and the Koch plate culture method.

Working with the D'yakonov apparatus, we were unable to obtain results which would satisfy us. For this reason only results of microbial contamination of air obtained with Krotov apparatus and the Koch plate culture are compared.

For the plate method test we used metal cans 10 cm in diameter and 20 cm in height and plates 9 cm in diameter. Glucose agar is poured into the latter. The volume of air in the can over the surface of the medium equalled to 1.5 liters.

We tested 3-5 cans at a time, i.e., we tested the microbe content of 4.5-7.5 liters of air.

Cans with the sterile air were opened in the room where air was to be sampled. The covered plates containing medium were placed in the cans on special supports. With several up-down movements the sterile air in the can was replaced with the air to be tested. The plate, containing medium, was uncovered and left at the bottom of the can. The can was closed and placed into an incubator for 48 hours. The bacteria in the column of air over the plate settled on the surface of the agar, grew and formed colonies, which we counted. Then we calculated the bacterial content in 1 m^3 of air.

In the same room and simultaneously with the collection of samples in cans we inoculated the air on the same medium with the Krotov apparatus.

The testing of the air was carried out in classrooms of schools and universities. The test results are presented in Table 1.

Table 1. Comparative data on the number of bacteria in 1 m^3 of air in a school.

| Class-room numbers | Air sampling | | | |
|--------------------|-----------------------|---------------|----------------|---------------|
| | with Krotov apparatus | | plate method | |
| | before classes | after classes | before classes | after classes |
| 1 | — | 2400 | 5000 | 15000 |
| 2 | 1933 | 4400 | 9000 | 19000 |
| 3 | 2266 | 4600 | 7000 | 28000 |
| 4 | 2100 | 8467 | 21000 | 46000 |
| 5 | 1167 | 7100 | 17000 | 133000 |
| 6 | 1900 | 5467 | 6600 | 13000 |
| 7 | 1922 | 2930 | 5730 | 19100 |
| 8 | 2233 | 10700 | 13000 | 55000 |
| 9 | 3500 | 6967 | 18600 | 36600 |

As is evident from the table, in the seeding of air with Krotov apparatus, a considerable number of bacteria are not caught. The average index of the number of bacteria in the air of the school during the testing with Krotov apparatus before classes is 5.3 times lower, and that after classes is 7 times lower, than that obtained with plate method.

Evidently, the air is not brought into total contact with the surface of the nutrient medium in the plate during its flow through the slit in the Krotov apparatus. As a result many bacteria, bypassing the medium, are ejected along with the outgoing air. To check this assumption, we placed opened plates containing agar in the path of the outgoing air and then counted the number of colonies. Subsequently by conversion we calculated the number of bacteria remaining in 1 m³ of air.

These data are presented in Table 2. Also, in this table the data are compared with the number of bacteria in 1 m³ according to the readings of Krotov apparatus with corrections from the plate culture method.

Table 2. Computation of the quantity of microorganisms in 1 m³ of air with different methods of air seeding.

| Sample number | Plate culture method | Krotov apparatus | | |
|---------------|----------------------|------------------|-----------------------|-------|
| | | Base plate count | Count in outgoing air | Total |
| 1 | 21000 | 2100 | 18000 | 20100 |
| 2 | 7000 | 4600 | 4000 | 8600 |
| 3 | 30230 | 2100 | 31160 | 33260 |
| 4 | 31020 | 2000 | 28000 | 30000 |
| 5 | 19700 | 1900 | 20000 | 21900 |

The presented data attest to the fact that Krotov apparatus does not trap a very large quantity of microorganisms.

In order to determine the effect of the rate of air flow through the apparatus on the test results, we carried out analogous studies with air flow at different rates (30, 20, and 10 liters per minute). The obtained results are presented in Table 3.

The data indicate that a reduced rate of air flow through the Krotov apparatus lowers the number of microorganisms settling on the nutrient medium even more.

Table 3. Computation of the quantity of microorganisms in 1 m³ of air with the Krotov apparatus with different rates of air flow.

| Test number | Rate of air flow in liters per minute | Base plate count | Count in outgoing air | Total |
|-------------|---------------------------------------|------------------|-----------------------|-------|
| 1 | 30 | 4880 | 10560 | 15540 |
| 2 | 20 | 3570 | 6070 | 9500 |
| 3 | 10 | 10900 | 32000 | 42900 |
| 4 | 30 | 7400 | 2600 | 10000 |
| 5 | 20 | 5880 | 6000 | 11880 |
| 6 | 10 | 6000 | 8000 | 14000 |
| 7 | 30 | 2100 | 10700 | 12800 |
| 8 | 20 | 1800 | 21000 | 22800 |
| 9 | 10 | 2900 | 9000 | 9300 |

It is obvious that the use of Krotov apparatus for seeding the air does not give satisfactory results. It is necessary to use the can-plate culture method, which gives better results.

UNCLASSIFIED

Security Classification

| DOCUMENT CONTROL DATA - R & D | | |
|--|---|---|
| (Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified) | | |
| 1. ORIGINATING ACTIVITY (Corporate author) Foreign Technology Division Air Force Systems Command U. S. Air Force | | 2a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED |
| | | 2b. GROUP |
| 3. REPORT TITLE COMPARATIVE CHARACTERISTIC OF SOME OF THE DEVICES USED TO DETERMINE THE MICROBIAL CONTAMINATION OF THE AIR | | |
| 4. DESCRIPTIVE NOTES (Type of report and inclusive dates) Translation | | |
| 5. AUTHOR(S) (First name, middle initial, last name) Ishchenko, G. N. , Khamrakulova, K. and Samigullin, R. | | |
| 6. REPORT DATE June 1962 | 7a. TOTAL NO. OF PAGES 4 | 7b. NO. OF REFS |
| 8a. CONTRACT OR GRANT NO. | 8b. ORIGINATOR'S REPORT NUMBER(S) FTD-HT-23-144-70 | |
| b. PROJECT NO. 6030024 | 8c. OTHER REPORT NO(S) (Any other numbers that may be assigned this report) | |
| c. | | |
| d. DIA Task No. T69-03-06 | | |
| 10. DISTRIBUTION STATEMENT Distribution of this document is unlimited. It may be released to the Clearinghouse, Department of Commerce, for sale to the general public. | | |
| 11. SUPPLEMENTARY NOTES | | 12. SPONSORING MILITARY ACTIVITY Foreign Technology Division Wright-Patterson AFB, Ohio |
| 13. ABSTRACT This article deals with a comparative study between Krotov apparatus and plate culture methods of determining microbial contamination of air. The air tested was that of schools and universities before and after classes. The results have been tabulated and comparison made. | | |

DD FORM 1473
1 NOV 60

UNCLASSIFIED

Security Classification

Security Classification

UNCLASSIFIED
Security Classification

Security Classification